

COMPOSITION

Expilin Tablet: contains Each tablet Chlorpheniramine Maleate BP 4 mg.

Expilin Syrup: Each 5 ml syrup contains Chlorpheniramine Maleate BP 2 mg.

PHARMACOLOGY

Expilin (Chlorpheniramine Maleate) is an alkylamine antihistamine. It is one of the most potent H, blocking agents and is generally effective in relatively low doses. Expilin is not so prone to produce drowsiness, readily absorbed from the gastro-intestinal tract, metabolised in the liver and excreted usually mainly as metabolised in the urine.

INDICATION

Expilin is mainly indicated in allergic conditions including urticaria, sensitivity reactions, angioneurotic oedema, seasonal hay fever, vasomotor rhinitis, cough, common cold, motion sickness.

DOSAGE & ADMINISTRATION

: 4 mg 3 - 4 times daily. Adults

Children: Up to 1 (one) year: 1 mg twice daily.

1 - 5 years : 1 mg 3 - 4 times daily. 6 - 12 years : 2 mg 3 - 4 times daily.

or as directed by the physician.

SIDE EFFECT

Drowsiness, dizziness, headache, psychomotor impairment, urinary retention, dry mouth, blurred vision and gastro-intestinal disturbances, paradoxical stimulation may rarely occur, especially in high dosage or in children.

CONTRAINDICATION & PRECAUTION

There is no definite contraindication to therapy. It should be used with caution in epilepsy, prostatic hypertrophy, glaucoma and hepatic disease.

Warning: The ability to drive or operate machinery may be impaired.

Store below 30°C. Protect from light. Keep out of reach of children.

PACKING

Expilin Tablet: Each box contains 10 x 50 tablet in

blister pack.

Expilin Syrup: Each bottle contain 100 ml syrup.

